

EVALUATION OF USE OF BIOGAS PLANT DIGESTATE AS FERTILIZER IN ALFALFA AND WINTER WHEAT

Milan Koszel¹, Artur Przywara¹, Francesco Santoro², Alexandros Sotirios Anifantis²

¹University of Life Sciences in Lublin, Poland; ²University of Bari Aldo Moro, Italy
milan.koszel@up.lublin.pl, artur.przywara@up.lublin.pl, francesco.santoro@uniba.it,
alexandrossotirios.anifantis@uniba.it

Abstract. Anaerobic decomposition of plant residues from which biogas is produced, generates very large amounts of digestate. Due to its physicochemical properties post-digestion liquid can be used as a fertilizer. Post-digestion liquid was used in the field cultivation of fodder alfalfa and winter wheat. The content of macroelements and the content of protein in the grains of winter wheat fertilized with digestate were on the same levels as in the grains of wheat fertilized with mineral fertilizers. The analysis showed a similar content of macroelements in alfalfa leaves fertilized both with post-digestion liquid and mineral fertilizers as well. Fertilizing fields with digestate brings destruction of possible pathogens. Digestate utilization as a fertilizer brings tangible benefits in agricultural production, but it is also a product, the application of which can reduce the negative effects of mineral fertilization and contribute to development of sustainable agriculture. The study has shown that digestate can be used as a fertilizer.

Keywords: digestate, fertilizer, alfalfa, winter wheat.

Introduction

Reduction of the amount of waste can be achieved by means of fermentation processes. Especially the process of anaerobic fermentation has enormous potential for the production of biofuels and bioproducts. Creating added value of bioproducts together with bioenergy opens up a number of opportunities of economic potential growth [1]. Anaerobic digestion of heterogenous waste, such as organic waste and other types of municipal waste, is a promising development of energy bioproduction from organic waste in the world [2] that could be used in several agricultural processes such as olive oil extraction [3-5], almonds postharvest process [6], packaging process [7].

Energy can be obtained in several sustainable ways, such as hydrogen systems coupled with renewable sources[8], geothermal energy, photovoltaic and solar thermal systems [9-11], boilers [12], microturbines [13] or from organic or municipal waste through the process of fermentation. This is a universally popular utilization method. During anaerobic digestion biogas and post-digestion liquid are produced, which can be used instead of mineral fertilizers [14-18]. Methane fermentation is a natural process – it is conducted in anaerobic conditions with the participation of microorganisms: acidogenic, acetogenic and methanogenic bacteria. In the process of methanogenesis most frequently there is decomposed approximately 30-60 % of organic matter introduced into the chamber together with the substrates [19]. Anaerobic digestion has long been used to produce biogas from organic residues, such as sewage sludge, agricultural and industrial by-products. Anaerobic decomposition converts organic waste into two products: biogas and fermented biomass (digestate). This biological process does not only produce renewable energy (biogas), but it also reduces the emission of greenhouse gases [20-22]. Particular care has to be taken in this biological process because of its intrinsic environmental and safety dangerousness like in other agricultural processes [23-25]. Anaerobic decomposition is a proven technology of biological waste processing; it entails degradation and stabilization of organic materials in anaerobic conditions by bacteria and leads to the production of biogas. The process is becoming a popular technology because of its ability to produce renewable energy from a wide variety of organic wastes from municipal, agricultural and industrial sources, while allowing nutrient recovery through application of the digestate to land. Post-digestion liquid is produced as well, which can be used as a fertilizer in agriculture. It should not be discharged into sewers due to the hazard of bacterial infestation [26].

Fermented biomass has the same or higher agricultural value than liquid manure since it contains more mineral components (including nitrogen) and less organic matter [20; 27; 28]. Digestate also has undesirable properties, such as: smell, viscosity, considerable humidity and high content of fatty acids, which are phytotoxic, this is why digestate can be pathogenic [20; 29].

The use of post-digestion liquid as a fertilizer brings substantial benefits for agriculture; the possibility of using fermented biomass as a fertilizer contributes to improved soil fertility and higher

crop yields. The utilization of post-digestion liquid as a fertilizer leads to the reduction of the use of mineral fertilizers [19; 30]. In literature [31-36] it is reported that digestate does not contain any heavy metals, and consequently can be used as a mineral fertilizer, which is an important step towards better environmental protection [37-40].

The aim of the work is to present the possibility of using a post-digestion liquid instead of a mineral fertilizer in field cultivation.

Materials and methods

Digestate obtained from the biogas plant in Piaski (Lubelskie Province) was applied on experimental fields for alfalfa and winter wheat. There were fodder alfalfa of Kometa variety and winter wheat of Zyta variety sown. For the sake of comparison, all the plants mentioned above were sown and fertilized with mineral fertilizers as well. Cultivation fields are located in the experimental farm of the University of Life Sciences in Lublin, in Czesławice Commune, Lubelskie Province. The area of each field was 75 m²: winter wheat in the first decade of September, 2016 and alfalfa, a perennial, in April, 2015 [41]. In the third year of harvesting (2017) alfalfa was harvested one time. The harvested plants were examined to determine the content of macroelements. Post-digestion liquid was used in the amount of 270 l per 75 m². On the field fertilized with mineral fertilizers, for alfalfa sowing, there were used: nitrogen – 20 kg·ha⁻¹, phosphorus – 60 kg·ha⁻¹, potassium – 80 kg·ha⁻¹ [41]. Then, on the field fertilized with mineral fertilizers, for winter wheat sowing, there were used: nitrogen – 140 kg·ha⁻¹ (first dose: pre-sowing and for spring pre-sowing cultivation, second dose in the period of shooting, third dose during earing), phosphorus – 62 kg·ha⁻¹, potassium – 81 kg·ha⁻¹.

Laboratory tests were performed at the District Chemical-Agricultural Station in Lublin and in the Central Agro-ecological Laboratory at the University of Life Sciences in Lublin.

Results and discussion

The post-digestion liquid was tested for the content of macroelements and heavy metals (Table 1). The pH of the digestate used for alfalfa and winter wheat cultivation was 8.39. The digestate sample contained no heavy metals. Moreover, the digestate contains substantial amounts of macroelements, therefore it can be used as a fertilizer. This is confirmed by the investigation conducted last year [30].

Soil samples were tested before and after the use of digestate and after the harvests of alfalfa and winter wheat. The results of the analyses are presented in Tables 2 and 3. The tests of the soil for alfalfa cultivation showed a very slight increase in pH reaction, from 7.46 to 7.50, then it dropped to 7.34 after the wheat harvest. A similar tendency was observed while testing the soil for winter wheat cultivation. The pH reaction increased from 7.59 to 7.60, and after the harvest its value dropped to 7.49. After winter wheat harvest the value of pH reaction was observed to fall to 7.49, which is a basic reaction. After the harvest of winter rape pH reaction value fell to 7.34, which is a neutral reaction.

Table 1

**Comparison of macroelements and heavy metals
in the digestate used for the field crops**

Examined feature	Content in digestate
Phosphorus, g·l ⁻¹	0.18
Potassium, g·l ⁻¹	6.12
Calcium, g·l ⁻¹	0.45
Magnesium, g·l ⁻¹	0.18
Cadmium, mg·l ⁻¹	0.40
Lead, mg·l ⁻¹	0.41
Nickel, mg·l ⁻¹	0.39
Chromium, mg·l ⁻¹	0.42
Copper, mg·l ⁻¹	0.52
Zink, mg·l ⁻¹	2.08
Manganese, mg·l ⁻¹	2.39
Iron, mg·l ⁻¹	74.66

Table 2

Tests for pH and macroelement content in the soil for alfalfa cultivation

Examined feature	Before digestate application	After digestate application	After harvest
Reaction, pH	7.46	7.50	7.34
Phosphorus, mg per 100 g of soil	34.19	45.02	40.58
Potassium, mg per 100 g of soil	8.30	13.89	10.55
Magnesium, mg per 100g of soil	14.49	18.23	15.01

Table 3

Tests for pH and macroelement content in the soil for winter wheat cultivation

Examined feature	Before digestate application	After digestate application	After harvest
Reaction, pH	7.59	7.60	7.49
Phosphorus, mg per 100 g of soil	49.48	55.87	52.41
Potassium, mg per 100 g of soil	16.01	17.93	16.78
Magnesium, mg per 100g of soil	13.28	18.68	17.05

In alfalfa cultivation, after digestate application, the content of phosphorus rose by 10.83 mg per 100 g of soil, potassium by 5.59 mg per 100 g of soil, and magnesium by 3.74 mg per 100 g of soil. As it was in the case of wheat cultivation, also in after alfalfa harvest there was observed decrease in the content of macroelements. In winter wheat cultivation the content of phosphorus rose by 6.39 mg per 100 g of soil, potassium by 1.92 mg per 100 g of soil and magnesium by 5.4 mg per 100 g of soil. After winter wheat harvest there was observed decrease in the content of macroelements. The decrease in the macroelement content after the harvests is connected with good absorption of macroelements by plants.

Table 4 shows changes in the content of macroelements in alfalfa leaves from the sixth harvest.

The highest rise was observed in the content of nitrogen, 0.11 p.p. The relative percent differences were: nitrogen – 3.15 %, phosphorus – 11.76 %, potassium – 1.00 %, calcium – 7.74 %, magnesium – 23.53 %.

Table 4

Content of macroelements in alfalfa leaves from the sixth harvest

Examined feature	Alfalfa sown on the soil fertilized with mineral fertilizers	Alfalfa sown on the soil fertilized with digestate
Nitrogen, %	3.49	3.60
Phosphorus, %	0.17	0.19
Potassium, %	1.99	2.01
Calcium, %	1.46	1.50
Magnesium, %	0.17	0.21

Winter wheat grains were collected from the fields fertilized with mineral fertilizers and digestate. The moisture of the grains collected from the field fertilized with mineral fertilizers was 12.9 %, and from the field fertilized with digestate – 13.4 %.

The elementary feature of wheat grains, which determines their value in use, is protein content. The protein content in the winter wheat grains collected from the field fertilized with mineral fertilizers was 10.84 %, and from the field fertilized with digestate 11.01 %. The relative percent difference for the protein content in the wheat grains is 1.56 %. The changes in the content of macroelements in winter wheat grains are presented in Table 5.

There was a slight percentage increase in the content of macroelements. The relative percent differences were: nitrogen – 3.93 %, phosphorus – 13.33 %, potassium – 9.26 %, calcium – 15.38 %, magnesium – 9.52 %.

Table 5

Content of macroelements in winter wheat grains.

Examined feature	Winter wheat sown on the soil fertilized with mineral fertilizers	Winter wheat sown on the soil fertilized with digestate
Nitrogen, %	1.78	1.85
Phosphorus, %	0.45	0.51
Potassium, %	0.54	0.59
Calcium, %	0.13	0.15
Magnesium, %	0.21	0.23

Conclusions

The examination of soil samples before and after digestate application showed increase in the content of macroelements in the soil, which implies a good fertilizing value of digestate. The analysis of the test results revealed a slight percentage increase in the content of particular macroelements in alfalfa and winter wheat. The examination of winter wheat grains from the soil fertilized with digestate also revealed a rise in the protein content as compared to the winter wheat grains from the field fertilized with mineral fertilizers. Fertilizing fields with digestate brings destruction of possible pathogens. Digestate utilization as a fertilizer brings tangible benefits in agricultural production, but it is also a product, the application of which can reduce the negative effects of mineral fertilization and contribute to development of sustainable agriculture.

References

- [1] Escamilla-Alvarado C., Poggi-Varaldo H. M., Ponce-Noyola M. T. Use of organic waste for the production of added-value holocellulases with *Cellulomonas flavigena* PR-22 and *Trichoderma reesei* MCG 80. *Waste Management & Research*, Vol. 31 (8), 2013, pp. 849-858.
- [2] Bayard R., Gonzalez-Ramirez L., Guendouz J., Benbelkacem H., Buffière P., Gourdon R. Statistical Analysis to Correlate Bio-physical and Chemical Characteristics of Organic Wastes and Digestates to Their Anaerobic Biodegradability. *Waste Biomass Valor*, 6, 2015, pp. 759-769.
- [3] Bianchi B., Tamborrino A., Santoro F. Assessment of the energy and separation efficiency of the decanter centrifuge with regulation capability of oil water ring in the industrial process line using a continuous method. *J Agr Eng* 2013;44(2s), 278-82. DOI: 10.4081/jae.2013.298
- [4] Bianchi B., Catalano P. Power consumption and quality features in the hammer crushing olive oil production. *GRACIAS Y ACETAS*, Vol. 47(3), 1996, pp. 136-141.
- [5] Caponio F., Squeo G., Brunetti L., Pasqualone A., Summo C., Paradiso VM., Catalano P., Bianchi B. Influence of the feed pipe position of an industrial scale two-phase decanter on extraction efficiency and chemical-sensory characteristics of virgin olive oil. *Journal of Food Engineering*, vol. 205, 2018, pp. 34-46. DOI: 10.1016/j.jfoodeng.2017.02.019
- [6] Pascuzzi S., Santoro F. Analysis of the almond harvesting and hulling mechanization process: A case study. *Agriculture (Switzerland)*, 7(12), 2017, 100. DOI: 10.3390/agriculture7120100
- [7] Manetto G., Cerruto E., Pascuzzi S., Santoro F. Improvements in citrus packing lines to reduce the mechanical damage to fruit. *Chem. Eng. Trans.* 58, 2017, pp. 391-396. DOI: 10.3303/CET1758066
- [8] Anifantis A. S., Colantoni A., Pascuzzi S., Santoro F. Photovoltaic and hydrogen plant integrated with a gas heat pump for greenhouse heating: A mathematical study. *Sustainability (Switzerland)*, vol. 10(2), 2018, pp. 1-12. DOI: /10.3390/su10020378
- [9] Anifantis A.S., Pascuzzi S., Scarascia Mugnozza G. Geothermal source heat pump performance for a greenhouse heating system: An experimental study. *J. Agric. Eng.*, vol.47, 2016, pp. 164-170. DOI: 10.4081/jae.2016.544
- [10] Anifantis A. S., Colantoni A., Pascuzzi S. Thermal energy assessment of a small scale photovoltaic, hydrogen and geothermal stand-alone system for greenhouse heating. *Renewable Energy*, vol. 103, 2017, pp. 115-127. DOI: 10.1016/j.renene.2016.11.031

- [11] Anifantis A. S. Performance assessment of photovoltaic, ground source heat pump and hydrogen heat generator in a stand-alone systems for greenhouse heating. *Chemical Engineering Transactions*, vol. 58, 2017, pp. 511-516. DOI: 10.3303/CET1758086
- [12] Bianchi B., Giametta F., La Fianza G., Gentile A. Catalano P. Microclimate measuring and fluid dynamic simulation in an industrial boiler house: testing of an experimental ventilation system. *Veterinaria Italiana*, Vol. 51 (2), 2015, pp. 85-92.
- [13] Perone C., Catalano F., Tamborrino A., Giametta F., Bianchi B., Ayr U. Study and Analysis of a Cogeneration System with Microturbines in a Food Farming of Dry Pasta. *Chem Engineer Trans*, vol.58, 2017, pp.499-504.
- [14] Comparetti A., Febo P., Greco C., Orlando S., Navickas K., Venslauskas K. Sicilian potential biogas production. *Journal of Agricultural Engineering*; vol. XLIV(s2):e103, 2013, pp. 522–525.
- [15] Bolzonella, D., Pavan, P., Mace, S., & Cecchi, F. Dry anaerobic digestion of differently sorted organic municipal solid waste: a full-scale experience. *Water Sci Tech*, vol.53(8),2006,pp.23-32.
- [16] Comparetti, A., Greco, C., Navickas, K., & Venslauskas, K. Evaluation of potential biogas production in Sicily. *Engineering for Rural Development*, vol. 11, 2012, pp. 555-559.
- [17] Sahlström, L. A review of survival of pathogenic bacteria in organic waste used in biogas plants. *Bioresource Technology*, vol. 87(2), 2003, pp. 161-166.
- [18] Weiland P. Biogas production: Current state and perspectives. *Applied Microbiology and Biotechnology*, vol. 85(4), 2010, pp. 849-860. DOI: 10.1007/s00253-009-2246-7.
- [19] Kowalczyk-Juśko A., Szymańska M. Poferment nawozem dla rolnictwa. *FnRRPR*, VA, 2015.
- [20] Bustamante M. A., Moral R., Bonmatí A., Palatsí J., Solé-Mauri F., Bernal M. P. Integrated Waste Management Combining Anaerobic and Aerobic Treatment: A Case Study. *Waste Biomass Valor*, 5, 2014. pp. 481-490.
- [21] Teglia C., Tremier A., Martel J.-L. Characterization of solid digestates: Part 1, review of existing indicators to assess solid digestates agricultural use. *Waste and Biomass Valorization*, vol. 2 (1), 2011, pp. 43-58. Cited 32 times. DOI: 10.1007/s12649-010-9051-5.
- [22] Holm-Nielsen J.B., Al Seadi T., Oleskowicz-Popiel P. The future of anaerobic digestion and biogas utilization. *Bioresource Technology*, vol. 100 (22), 2009, pp. 5478-5484.
- [23] Pascuzzi S., Santoro F. Analysis of possible noise reduction arrangements inside olive oil mills: A case study. *Agriculture (Swit.)*, 7(10), 2017, 88. DOI: 10.3390/agriculture7100088
- [24] Pascuzzi S., Santoro F. Evaluation of farmers' OSH hazard in operation nearby mobile telephone radio base stations. 16th International Scientific Conference "Engineering for rural development" Proceedings, Volume 16. Jelgava, Latvia, May 24-26, 2017, pp. 748-755 – ISSN: 1691-5976, DOI: 10.22616/ERDev2017.16.N151
- [25] Pascuzzi S., Santoro F. Exposure of farm workers to electromagnetic radiation from cellular network radio base stations situated on rural agricultural land. *International Journal of Occupational Safety and Ergonomics*, 21(3), 2015, pp. 351-358. DOI: 10.1080/10803548.2015.1081774
- [26] Beneragama N., Moriya Y., Yamashiro T., Iwasaki M., Lateef S. A., Ying C., Umetsu K. The survival of cefazolin-resistant bacteria in mesophilic co-digestation of dairy manure and waste milk. *Waste Management & Research*, 31 (8), 2013, pp. 843-848.
- [27] Abdullahi Y.A., Akunna J.C., White N.A., Hallett P.D., Wheatley R. Investigating the effects of anaerobic and aerobic post-treatment on quality and stability of organic fraction of municipal solid waste as soil amendment. *Bioresource Technology*, vol. 99 (18), 2008, pp. 8631-8636.
- [28] Iglesias J.E., Pérez G.V. Determination of maturity indices for city refuse composts. *Agriculture, Ecosystems and Environment*, vol. 38 (4), 1992, pp. 331-343.
- [29] Farrell M., Jones D.L. Critical evaluation of municipal solid waste composting and potential compost markets. *Bioresource Technology*, vol. 100 (19), 2009, pp. 4301-4310.
- [30] Koszel M., Kocira A., Lorencowicz E. The evaluation of the use of biogas plant digestate as a fertilizer in alfalfa and spring wheat cultivation. *Fresen Environ Bull*, 25(8), 2016, pp. 3258-3264.
- [31] Odlare M., Pell M., Svensson K. Changes in soil chemical and microbiological properties during 4 years of application of various organic residues. *Waste Management*, 28, 2008, pp. 1246–1253.
- [32] Montemurro F., Vitti C., Diacono M., Canali S., Tittarelli F., Ferri D. A three-year field anaerobic digestates application: effects on fodder crops performance and soil properties. *Fresenius Environmental Bulletin*. Volume 19 – No 9b, 2010, pp. 2087-2093.

- [33] Salminen E., Rintala J., Härkönen J., Kuitunen M., Högmander H., Oikari A. Anaerobically digested poultry slaughterhouse wastes as fertiliser in agriculture. *Bioresource Technology*, vol. 78 (1), 2001, pp. 81-88.
- [34] Melamane X.L., Tandlich R., Burgess J.E. Submerged membrane bioreactor and secondary digestion for the treatment of wine distillery wastewater. Part I: Raw wine distillery wastewater digestion. *Fresenius Environmental Bulletin*, vol. 16 (2), 2007, pp. 154-161.
- [35] Moletta R. Winery and distillery wastewater treatment by anaerobic digestion. *Water Science and Technology*, vol. 51 (1), 2005, pp. 137-144.
- [36] Bustamante M.A., Paredes C., Moral R., Moreno-Caselles J., Pérez-Espinosa A., Pérez-Murcia M.D. Uses of winery and distillery effluents in agriculture: Characterisation of nutrient and hazardous components. *Water Science and Technology*, 51 (1), 2005, pp. 145-151.
- [37] Montemurro F., Canali S., Convertini G., Ferri D., Tittarelli F., Vitti C. Anaerobic digestates application on fodder crops: Effects on plant and soil. *Agrochimica*, 52 (5), 2008, pp. 297-312.
- [38] Tambone F., Genevini P., D'Imporzano G., Adani F. Assessing amendment properties of digestate by studying the organic matter composition and the degree of biological stability during the anaerobic digestion of the organic fraction of MSW. *Bioresource Technol.*, vol. 100(12), 2009, pp. 3140-3142.
- [39] Canali S., Trinchera A., Intrigliolo F., Pompili L., Nisini L., Mocali S., Torrisi B. Effect of long term addition of composts and poultry manure on soil quality of citrus orchards in Southern Italy. *Biology and Fertility of Soils*, vol. 40 (3), 2004, pp. 206-210.
- [40] Douglas J.T., Aitken M.N., Smith C.A. Effects of five non-agricultural organic wastes on soil composition, and on the yield and nitrogen recovery of Italian ryegrass. *Soil Use and Management*, vol. 19(2), 2003, pp. 135-138. DOI: 10.1079/SUM2003180.
- [41] Koszel M, Lorencowicz E. Agricultural use of biogas digestate as a replacement fertilizers. *Agriculture and Agricultural Science Procedia* 7, 2015, pp. 119-124.